

Shiraz University of Medical Sciences

Shahid Dastgeyb hospital



## **Functional Endoscopic Sinus surgery Post- Operative care Instruction Sheet**

producer: A. Rezaiefard

Corroborant: Dr. Tayebbeh Kazemi

2023

**CALL THE OFFICE at  
+989355553898 if you  
have any questions**

**Reference:  
[www.clevelandclinic/head  
andneck](http://www.clevelandclinic/headandneck)**

You may experience some discomfort postoperatively due to manipulation and inflammation. Take your pain medication as directed. Often extra-strength Tylenol is sufficient. You may wish to take medication for pain prior to your post-operative visits (particularly early on, when the nose is most sensitive). If the medication is sedating, be sure to have someone available to drive you. Do not hesitate to contact our office for any questions or for any possible emergency situation. If necessary, do not hesitate to go to the emergency department nearest to you for immediate care.

In general, you may resume other medications on the day after surgery. If you have any questions, please ask your physician.

Please call us immediately if you note fever (temperature above 38o C) clear watery drainage from your nose, changes in your vision, eye swelling, worsening headache and/or neck stiffness.

**What can I expect to experience during the post-op visits?**

You can expect some bleeding from your nose for several days after the surgery and again after each office debridement. When bleeding occurs down the front of your nose or into the back of your throat, you should tilt your head back while sitting up and breathe gently through your nose. If bleeding persists for an extended period of time, please call the office.

As the sinuses begin to clear themselves, you can expect to have some thick brown drainage from your nose. This is mucus and old blood and does not indicate an infection.

۴

Start your nasal irrigations on the day after surgery. You will be given instructions on how to perform the irrigations.

Do not blow your nose for two weeks after surgery.

(Occasionally, you will be given instructions not to blow your nose for longer periods of time.)

Do not bend, lift or strain for two weeks after surgery. These activities will promote bleeding from your nose. You should not participate in rigorous activity until healing is completed.

Do not suppress the need to cough or sneeze; instead, cough or sneeze with your mouth open.

Do not take aspirin and nonsteroidal anti-inflammatory drugs (ibuprofen, naproxen, Advil, Motrin, Aleve, others) for a full 7-10 days after surgery.

Take your prescribed antibiotics as directed by your physician.

If you are taking blood-thinning drugs (Coumadin, Plavix) on the recommendation of a physician, please ask your physician about the best time to resume these medications.

۳

**About Post-Operative Care Visits**

Post-operative office visits are an indispensable part of the surgery, since they help promote healing and prevent persistent or recurrent disease. During these visits, the surgical cavity is cleaned and inspected. Early scar tissue may be removed, and the medical treatment strategy will be adjusted.

Your first post-operative visit will occur typically within the first week after surgery, and you should expect 2 post-operative visits within the first 2-3 weeks after surgery.

Although complications from the manipulations performed during the post-operative visits are very rare, the theoretical risks are the same as the surgery itself. Consent for surgery includes consent for post-operative care, since the surgery and post-operative care are so closely related.

Failure to follow the postop care regimen can adversely affect the outcome of the surgery.

**Important Instructions for the Post-Operative**

۲